

Effect of hydrophobicity of post-tanning materials on microstructure and properties of crust leather


Wenjun Long¹, Liangqiong Peng¹, Xindi Lv¹, Yue Sun¹, Yue Yu^{1*}, Wenhua Zhang^{1,2*}

1. Key Laboratory of Leather Chemistry and Engineering of Ministry of Education, Sichuan University, Chengdu 610065, China


2. National Engineering Laboratory for Clean Technology of Leather Manufacture, Sichuan University, Chengdu 610065, China

Introduction


- ⊙ The hydrophobic post-tanning material enhances the fiber dispersion and mechanics of the crust leather.
- ⊙ The hydrophobic post-tanning material regulates the water types and fiber bundles radius of the crust leather.
- ⊙ The fiber content/fiber bundle radius ((1 - P)/R) is proportional to the tear strength of the crust leather.



Results




- ⊙ Water content of tanned leather during ethanol dehydration (a); Water content (b), tear strength (c), elongation at break (d), softness (e), tensile strength (f) of crust leather treated with amino silicone oil.




- ⊙ MIP curves of crust leather treated with different amino silicone oils ((a) Cr-leather; (b) TWLZ-leather; (c) TWS-leather); Average pore radius of crust leather (d); Tanning FB radius (e); Fitting of average pore size and FB radius of crust leather (f).


Results



- ⊙ Rehydrated curves of 12 kinds of crust leather ((a) Cr-, (b) TWLZ-, (c) TWS-leather).



- ⊙ Fitting of (1 - P)/R to the tear strength according to the crack-bridging model



- ⊙ SAXS curves of (a) Cr-leather, (b) TWLZ-leather and (c) TWS-leather.

Conclusions

- ⊙ Amino silicone oil can reduce the multi-layer water and side chain water content in the crust leather.
- ⊙ Amino silicone oil can regulate the multi-layer pore structure in leather and improve the D-spacing of fibrils, and promote the formation of smaller pores and better dispersed fiber bundles in the crust leather.
- ⊙ The linear correlation between tear strength and (1-P)/R validates FB as a bearing element for the mechanical properties of leather materials.

Acknowledgments

- ⊙ This project was funded by the National Natural Science Foundation of China (No. 22278278).